

CardiologyRounds®

www.cardiologyrounds.org

AS PRESENTED IN THE ROUNDS OF THE CARDIOVASCULAR DIVISION
OF BRIGHAM AND WOMEN'S HOSPITAL, BOSTON, MASSACHUSETTS

Fish Intake, Contaminants, and Human Health: Evaluating the Risks and the Benefits Part 2 – Health Risks and Optimal Intakes

By DARIUSH MOZAFFARIAN, MD, DRPH

This two-part report summarizes findings from a recent publication in the *Journal of the American Medical Association*.¹ Part 1, in the October issue of *Cardiology Rounds*, discussed the potential health benefits of fish and fish oil intake. Part 2, in this issue, discusses the potential health risks and describes optimal intakes for different populations. The tables and figures in this report are adapted from *JAMA*, October 18, 2006; 296:1886-99, © 2006, American Medical Association, all rights reserved.

Health effects of fish intake – controversy and confusion

Since low rates of coronary heart disease (CHD) death were observed in Greenland Eskimos consuming high quantities of seafood,² evidence from animal-experimental, observational, and clinical research has further supported a protective effect of fish consumption on risk of CHD death. Two marine n-3 polyunsaturated fatty acids (n-3 PUFAs) – eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) – have been identified as the likely active constituents.³⁻²¹ On the other hand, there is growing concern over the potentially harmful effects of mercury, dioxins, and polychlorinated biphenyls (PCBs) that are present in some fish species.²²⁻²⁹ As a result, there are contradictory messages on the risks and benefits of fish intake, causing controversy and confusion over the health effects of fish consumption.^{30,31}

Health risks of mercury

Mercury is emitted into the atmosphere from several sources, including volcanoes, coal-fired electric power plants, gold mining, institutional boilers, chlorine production, and waste incineration. From the atmosphere, mercury cycles from rainwater into lakes and oceans, where it is converted by the action of microorganisms into organic methylmercury. Compared with inorganic or elemental mercury, methylmercury bioaccumulates in aquatic food chains and has greater potential toxicity than inorganic mercury.^{22,23,25} For example, inorganic mercury is poorly absorbed in the gastrointestinal tract following consumption, and elemental mercury does not readily cross tissue barriers, while methylmercury is well absorbed and actively transported into tissues by a widely distributed carrier protein.²²

The concentration of methylmercury in any given fish species depends on the degree of local environmental contamination and on the predatory nature and lifespan of the species. Longer-living, larger predators, such as swordfish or shark, have higher tissue concentrations, while shorter-lived or smaller species, such as salmon or shellfish, have very low concentrations (Table 1). Methylmercury content of fish is not significantly altered by preparation methods.²²

Following uptake into tissues, mercury can bind to the sulfhydryl groups of enzymes, ion channels, and receptors, which can inhibit important antioxidant systems and increase the production of reactive oxygen species and free radicals.^{22,24} Health effects of very high doses of mercury exposure following occupational or industrial accidents are well-documented and include paresthesias, ataxia, and sensory abnormalities in adults, and delayed cognitive and neuromuscular development in children following *in utero* exposure.^{22,32} However, the health effects of chronic low level mercury exposure – i.e., that seen with typical fish consumption – are less well-established. This evidence is reviewed herein.

Early neurodevelopment

Methylmercury crosses the placenta, and exposure to the fetus is a function of maternal exposure.³³ Following very high gestational exposure, severe neurodevelopmental abnormalities can occur in children. This was seen, for example, following industrial accidents in Japan in the 1950s and Iraq in 1971, when



BRIGHAM AND
WOMEN'S HOSPITAL



HARVARD
MEDICAL SCHOOL
TEACHING AFFILIATE

Cardiovascular Division (Clinical)

Dale Adler, MD	Jane A. Leopold, MD
Christine Albert, MD	Eldrin Lewis, MD
Michelle Albert, MD	James Liao, MD
Elliott Antman, MD	Peter Libby, MD (Division Chief)
Kenneth Baughman, MD	Leonard Lilly, MD
Joshua Beckman, MD	Bernard Lown, MD
Charles M. Blatt, MD	Laura Mauri, MD
Eugene Braunwald, MD	Thomas Michel, MD, PhD
Christopher Cannon, MD	David Morrow, MD
Ming Hui Chen, MD	Karen Moulton, MD
Alanna Coolong, MD	Gilbert Mudge, MD
Mark Creager, MD	Anju Nohria, MD
Akshay Desai, MD	Patrick O'Gara, MD
Elazer Edelman, MD, PhD	Marc A. Pfeffer, MD, PhD (Editor)
Andrew Eisenhauer, MD	Jorge Plutzky, MD
Laurence Epstein, MD	Jeffrey Popma, MD
David Faxon, MD	Shmuel Ravid, MD
Mark Feinberg, MD	Frederic Resnic, MD
Daniel Forman, MD	Paul Ridker, MD
Peter Ganz, MD	Thomas Rocco, MD
J. Michael Gaziano, MD	Maria Rupnick, MD, PhD
Thomas Gaziano, MD	Marc Sabatine, MD
Marie Gerhard-Herman, MD	Arthur Sasahara, MD
Robert Giugliano, MD	Benjamin Scirica, MD
Michael Givertz, MD	Christine Seidman, MD
Samuel Z. Goldhaber, MD	Andrew Selwyn, MD
Thomas B. Graboys, MD	Laurence Sloss, MD
Howard Hartley, MD	Piotr Sobieszcyk, MD
Carolyn Ho, MD	Regina Sohn, MD
John Jarcho, MD	Scott Solomon, MD
Paula Johnson, MD	Lynne Stevenson, MD
Scott Kinlay, MD	William Stevenson, MD
Jamil Kirdar, MD	Peter Stone, MD
James Kirshenbaum, MD	Michael Sweeney, MD
Bruce Koplan, MD	Usha Tedrow, MD
Raymond Kwong, MD	Stephen Wiviott, MD
Michael J. Landzberg, MD	Justina Wu, MD
Richard Lee, MD	

Brigham and Women's Hospital

Fax: (617) 732-5291 Website: www.heartdoc.org

The editorial content of *Cardiology Rounds* is determined solely by the Cardiovascular Division of Brigham and Women's Hospital. This publication is made possible by an educational grant.

**Cardiology Rounds is approved
by the Harvard Medical School
Department of Continuing Education
to offer continuing education credit**

Table 1: Levels of n-3 fatty acids and contaminants in commonly consumed fish, shellfish, and other foods

	EPA + DHA, mg per serving (serving size)	Selenium, µg/g (parts per million)	Mercury, µg/g (parts per million)	PCBs, ng/g (parts per billion)	Dioxins, TEQ pg/g (parts per trillion)
FDA action level	n/a	n/a	1.0	2000	none
Fish					
Anchovy	1165 (2 oz)	0.68	<0.05		0.35 (1997-98)
Catfish, Farmed	253 (5 oz)	0.15	<0.05	<50 (1997)	0.52 (1995-97)
Cod, Atlantic	284 (6.3 oz)	0.38	0.10		0.10 (1995-97)
Fish Burger, Fast Food	337 (2.2 oz)	0.17	<0.05	8 (2001)	0.06 (2001)
Fish Sticks, Frozen	193 (3.2 oz)	0.17	<0.05		0.04 (2001)
Golden Bass (Tilefish)	1,358 (5.3 oz)	0.52	1.45		
Halibut	740 (5.6 oz)	0.47	0.25		1.00 (1995-97)
Herring, Atlantic	1,712 (3 oz)	0.47	<0.05		0.97 (1995-98)
Mackerel, Atlantic	1,059 (3.1 oz)	0.52	0.05		0.60 (1997-98)
Mackerel, King	618 (5.4 oz)	0.47	0.73		
Salmon, Farmed†	4,504 (6 oz)	0.41	<0.05	25* (2001-03)	0.61 (2001-03)
Salmon, Wild†	1,774 (6 oz)	0.46	<0.05	1.8* (2002)	0.19 (2002)
Sardines	556 (2 oz)	0.53	<0.05	40 (2001-03)	0.31 (2001-03)
Shark	585 (3 oz)	0.34	0.99		
Swordfish	868 (3.7 oz)	0.62	0.98		
Trout	581 (2.2 oz)	0.15	0.07	11 (2002)	0.44‡ (2002)
Tuna, Light (Skipjack)†	228 (3 oz)	0.80	0.12	45 (2001)	0.02 (1995-98)
Tuna, White (Albacore)†	733 (3 oz)	0.66	0.35	100 (2001-03)	0.23 (2001-03)
Shellfish					
Crab	351 (3 oz)	0.40	0.09	6 (2002)	0.55‡ (2002)
Scallops	310 (3 oz)	0.28	<0.05		0.16 (1998)
Shrimp	267 (3 oz)	0.40	<0.05	1.1 (2002)	0.09‡ (2002)
Other foods					
Beef	0	0.19	0	22 (2001)	0.20 (1995-2001)
Butter	0	<0.05	0	70 (2001)	0.44 (1995-2001)
Cheese	0	0.22	0		0.45 (1995-2001)
Chicken	0	0.23	0	32 (2001)	0.11 (1995-2001)
Eggs	22 (1 egg)	0.23	0	19 (2001)	0.29 (1998-2001)
Milk	0	0.02	0		0.07 (1995-2001)

Average values shown (Mozaffarian and Rimm³), which may vary due to methodologic, geographic, temporal, and fish-to-fish differences. Values for PCBs and dioxins may overestimate current levels because contaminant levels in most foods, including fish species, are decreasing over time (eg, TEQs decreased by 33%-81% in meats and 66%-77% in salmon and tuna fish between 1995 and 2003); year of sampling is given in parenthesis.

* Values including the fish skin; levels may be lower in the edible portion.

† For the same specific species, there are minimal differences in nutritional or contaminant content of canned vs. fresh salmon or tuna. However, different species are typically canned vs. sold fresh. For salmon, differences between species are small compared with differences between farmed and wild salmon. For tuna, canned light (skipjack) tuna and fresh yellowfin/ahi tuna are more similar overall, while canned white (albacore) tuna and fresh bluefin tuna are more similar overall.

‡ Includes dioxin-like PCBs. TEQ = Toxic equivalence.

mothers consumed highly contaminated fish (mercury content = 10-30 ppm) or highly contaminated grain (maternal intake = 710-5700 µg/kg/d; maternal hair mercury = 18-598 ppm), respectively.^{22,32} More typical methylmercury exposures from fish consumption are far lower. For example, among U.S. women of childbearing age, the median (10th, 95th percentiles) levels of hair mercury were 0.19 ppm (0.04, 1.73) overall, and 0.34 ppm (0.09, 2.75) among women who consumed 3+ servings of fish per month.³⁴ These low exposure levels do not produce clinically detectable neurologic symptoms or signs in children. Several prospective studies have evaluated whether subclinical effects, detectable with specialized testing, might occur.³⁵⁻⁴³ In studies in the Faroe Islands,^{35,36} New Zealand,^{37,38} and Poland,³⁹ higher gestational mercury exposure was associated with lower scores on some neurologic tests (eg, finger tapping or naming tests), but not on most other neurologic tests. In contrast, in the Seychelles, higher gestational mercury exposure was associated with *higher* scores on some neurologic tests.^{40,41}


In a study of U.S. infants, maternal fish intake during gestation was associated with better visual recognition memory scores, while maternal hair mercury was associated with lower visual recognition memory scores,⁴² suggesting that overall fish consumption (which provides DHA, likely beneficial for neurodevelopment; see Part 1) and methylmercury exposure may have opposing effects. Among British children, gestational mercury exposure was not associated with neurodevelopmental scores; in

contrast, both maternal and infant fish intake were associated with better neurodevelopmental scores.⁴³ Several other studies have not seen consistent associations between gestational mercury exposure and neurologic test scores during childhood.⁴⁴

Direct comparisons of these different studies are challenging, due to variation in study designs (prospective vs. cross-sectional), methods for measuring mercury exposure, neurologic tests employed, timing of assessment (infancy vs. childhood), and statistical methods. Some analyses are also limited by incomplete adjustment for potential confounding risk factors, such as socioeconomic status, and multiple statistical testing (eg, 30+ neurologic parameters), increasing the possibility of chance associations. Randomized trials to test the effects of reducing low-level methylmercury exposure during gestation have not been performed. Nevertheless, because some associations with lower neurologic test scores are seen in some studies, and because clinical neurotoxicity is seen following much higher level accidental exposures, it is prudent to conclude that lower exposure levels may produce subclinical neurodevelopmental deficits.

Based on this, the Environmental Protection Agency published a focused Advisory for women of childbearing age, nursing mothers, and young children.⁴⁵ The lower 95% confidence limit at which gestational mercury exposure may produce abnormal subclinical neurologic test scores was determined to be 1 µg/kg/day, or ~500 µg/week for a 70 kg woman.³³ For additional safety, this was multiplied by a 10-fold uncertainty factor,

Figure 1: Studies of mercury exposure and risk of coronary heart disease (CHD). Relative risk (•) and 95% CIs (–) are shown comparing the highest to the lowest quantile of mercury exposure after adjustment for other risk factors. Combining results of studies, the relationship was not significant (RR= 1.12, 95% CI, 0.71-1.75, $p=0.62$), with significant heterogeneity between studies (p heterogeneity=0.008).



resulting in a reference dose, the allowable upper limit of daily intake, for methylmercury of 0.1 $\mu\text{g}/\text{kg}/\text{day}$ ($\sim 50 \mu\text{g}/\text{week}$ for a 70 kg woman).³³ Four fish species (shark, swordfish, king mackerel, tilefish) exceed this limit in a single serving (Table 1), so the Advisory recommended that women of childbearing age, nursing mothers, and young children avoid these specific species.⁴⁵ Notably, the Advisory also emphasized that women and children should consume a variety of other fish up to 2 servings/week (including up to 1 serving/week of albacore tuna) to receive the important health benefits.⁴⁵ It is also relevant that this Advisory was intended only for this specific population, because the importance of this conservative reference dose for health effects in adults is unclear.⁴⁶

Cardiovascular disease

The results of studies of mercury exposure and cardiovascular disease incidence in adults are conflicting (Figure 1),⁴⁷⁻⁵¹ providing inconclusive evidence for cardiovascular toxicity of mercury exposure. Of note, in the only 2 studies that observed positive associations between mercury exposure and cardiovascular risk, the *net* effect of fish consumption was still beneficial: compared to those not eating fish or n-3 PUFAs, individuals consuming fish or n-3 PUFAs still had lower risk of heart disease, but the size of this lower risk was somewhat less if their mercury exposure was higher.^{49,51,52} This indicates that the major public health question may not be whether consumption of mercury-containing fish increases cardiovascular risk, but whether consumption of such fish would decrease risk even *further* if mercury were not present. This is an important policy question: the potentially attenuated benefits of fish intake due to mercury content must be balanced against the costs and practicality of reducing mercury levels in the environment. However, this question should not obscure the strong evidence for the net cardiovascular benefits of fish consumption (see Part 1). This balance of benefit vs. risk would be most favorable for oily fish species which contain higher amounts of n-3 PUFAs (ie, most mercury-containing ocean fish), compared with lean fish from freshwater sources, which are generally lower in n-3 PUFAs.

Neurologic outcomes

Sensorimotor symptoms in adults, most commonly paresthesias, can be seen following very high methylmercury exposure from accidents^{22,32,53} or prolonged high intakes of

mercury-containing fish (e.g., 1-2 fish servings per day, including species high in mercury, for >10 years⁵⁴). Such symptoms are typically reversible when mercury exposure is reduced.

Lower levels of mercury exposure (that seen with typical fish intake) do not produce clinically detectable neurologic symptoms or signs in adults. Cross-sectional studies have evaluated whether chronic low level mercury exposure (assessed by hair or blood mercury levels) is associated with subclinical neurologic measures in adults. In studies among Amazon basin and Quebec Cree individuals, both positive and inverse associations were seen between mercury levels and some neurologic measures.⁵⁵⁻⁵⁷ These results were limited by multiple statistical testing (typically 20-30+ neurologic tests or participant subgroups were evaluated) and by possible inadequate adjustment for other potential risk factors. In a study among U.S. adults, mercury levels were associated with lower visual memory scores ($p=0.01$) but better motor and manual dexterity scores ($p=0.02$) among 20 different outcomes evaluated.⁵⁸ Among elderly Swedish adults, no associations were found between mercury levels and cognitive function.⁵⁹ Thus, these studies do not provide clear evidence that low-level methylmercury exposure affects subclinical neurologic outcomes in adults and, if so, what quantities or durations of exposure might be necessary. On the other hand, growing evidence suggests that fish consumption may favorably affect *clinical* neurologic outcomes in adults, including ischemic stroke,⁶⁰ cognitive decline and dementia,⁶¹ and depression and other neuropsychiatric disorders.^{62,63}


Health risks of dioxins and PCBs

Dioxins (commonly referring to dibenzodioxins and dibenzofurans) are organochlorine by-products of industrial processes involving chlorine, such as waste incineration, paper bleaching, pesticide production, and production of polyvinyl chloride plastics.²⁸ Regulatory and industry efforts have reduced dioxin emissions $>90\%$ since 1987.²⁸ PCBs are synthetic organochlorine compounds previously used in industrial and commercial processes.²⁹ Manufacture and processing of PCBs was prohibited in 1977.²⁹ Thus, environmental levels of these contaminants are steadily declining.^{28,64-67} Nevertheless, these contaminants persist for long periods in the environment, and PCBs and dioxins continue to be present in low concentrations in many foods (Table 1).

Cancer risks: Based upon animal experiments and some limited evidence in humans, PCBs and dioxins are carcinogenic.^{27,68} There are multiple different chemical structures (congeners) of dioxins and PCBs, each with different potential carcinogenicity. To calculate the potential toxicity of any food, the level of each congener in the food is multiplied by that congener's toxic equivalency factor (standardized against 2,3,7,8-tetrachloro-dibenzo-p-dioxin), and these values are summed to determine the Toxic Equivalence (TEQ), the measure of overall potential toxicity. In the U.S., dioxins comprise 72% and PCBs 28% of total TEQ exposure.⁶⁹ The major dietary sources of PCBs/dioxins are not fish and shellfish (only 9% of total TEQ), but beef, chicken, and pork (34% of total TEQ); dairy products (30% of total TEQ); and vegetables (22% of total TEQ).⁶⁹

Although the major dietary sources of PCB/dioxin exposure are meats, dairy products, and vegetables, much attention has been given to levels in fish (Table 1). When PCBs and dioxins were measured in farmed and wild salmon,^{70,71} levels were similar to several other foods (Table 1). Substantial amounts of n-3 PUFAs were also present in farmed and wild salmon: 4,504 and 1,774 mg EPA+DHA per 6-oz, respectively.⁷¹ A quantitative

Figure 2: Estimated cancer risks and CHD benefits of consumption of farmed and wild salmon, assuming regular intake to provide 1 g EPA+DHA per day over a 70-year lifetime. Based on animal-experimental data and limited studies in humans at much higher doses (and including a 10-fold safety factor), farmed and wild salmon consumption would result in 24 and 8 excess cancer deaths, respectively, per 100,000 individuals. Conversely, based on prospective observational studies and randomized clinical trials in humans, consumption of either farmed or wild salmon would result in 7,125 fewer CHD deaths.



risk-benefit analysis evaluated the potential cancer risks vs. CHD benefits, assuming regular consumption of farmed or wild salmon to provide 1000 mg/d of EPA+DHA over a 70-year lifetime (Figure 2).^{72,73} Such consumption would result in 24 and 8 excess cancer deaths per 100,000 lifetimes for farmed and wild salmon, respectively, but also 7,125 fewer CHD deaths per 100,000 lifetimes for either farmed or wild salmon.⁷² Thus, for both farmed and wild salmon, the potential benefits far exceeded the potential risks. When age-specific estimates were evaluated,¹ CHD benefits outweighed cancer risks at any age (from 25-34 to 85+ years) by 100-fold to 370-fold for farmed salmon and by 300-fold to >1000-fold for wild salmon.

It is important to note that the estimated CHD benefits are based upon prospective studies and randomized trials in humans (see Part 1), while the estimated cancer risks are based on animal-experimental data and limited studies in humans at high doses. The estimated cancer risks also include a ten-fold safety factor⁷³ – i.e., the potential risk may be as low as 2.4 and 0.8 cancer deaths per 100,000 lifetimes for farmed and wild salmon, respectively. The estimated cancer risks also assumed a lifetime consumption of salmon to provide 1000 mg/day of EPA+DHA, roughly corresponding to four 6-oz wild salmon every week for 70 years. However, reductions in CHD mortality are likely achieved with much lower intake: ~250 mg/d (see Part 1), or one 6-oz wild salmon serving/week. At this lower intake, CHD benefits would be largely unchanged (~7,125 fewer CHD deaths) while lifetime cancer risk would decrease ~75% (6 and 2 estimated deaths per 100,000 lifetimes for farmed and wild salmon, respectively, or 0.6 and 0.2 estimated deaths per 100,000 lifetimes not including the ten-fold safety factor). Consistent with these very low cancer risks, prospective studies in humans have seen little evidence for effects of fish intake on cancer risk.⁷⁴

Other risks: PCBs and dioxins may have non-cancer risks in adults, such as neurologic or immune system effects.²⁷⁻²⁹ On the other hand, consumption of fish may also result in other benefits, such as lowering risk of other cardiovascular outcomes (see Part 1), dementia,⁶¹ neuropsychiatric disorders,

^{62,63} and inflammatory disorders.^{75,76} To alter the estimates of risks vs. benefits meaningfully, such additional possible risks would have to exceed the additional possible benefits by more than 100-fold.

Exposure to PCBs and dioxins during gestation (but not infancy) has been associated with childhood neurodevelopmental deficits in several,⁷⁷⁻⁸³ though not all,^{84,85} studies. Because most exposure (>90%) is from meat, dairy, and vegetable sources,^{69,86} this concern is not specific to fish consumption, particularly since fish also contains DHA which has likely neurodevelopmental benefits. Nevertheless, because some freshwater fish may have higher levels of PCBs/dioxins, women of childbearing age who consume locally caught freshwater fish from highly contaminated inland sources or who frequently consume (1+ servings/day) commercially sold freshwater fish⁸⁶ should consult regional advisories.

Fish oil supplements: Levels of PCBs (0-450 ng/g,^{87,88}) and dioxins (0.2-11 TEQ pg/g^{89,90}) in fish oil are a function of the levels in the fish species from which the fish oil is derived. Given the small amounts of fish oil consumed (1-3 g/d), the total PCB/dioxin exposure from fish oil intake is extremely low. Fish oil capsules contain very little to no mercury.⁹¹

Summary of the evidence: optimal intakes

The optimal target intake of n-3 PUFAs may depend on both the population and the outcome of interest. As detailed in Part 1, based on the sum of evidence from prospective studies, clinical trials, and experiments, fish or fish oil intake reduces the risk of CHD death/sudden death. In the general population, 250 mg/d EPA+DHA is a reasonable target intake to reduce CHD mortality. Dietary n-3 PUFAs persist for weeks in tissue membranes,⁹² and so this amount can be converted to weekly intake of ~1500-2000 mg. This level of intake can be achieved by consuming one 3-oz serving/wk of farmed salmon, one 6-oz serving/wk of wild salmon or similar oily fish, or more frequent intake of less n-3 PUFA-rich fish (Table 1).

For individuals with established CHD, the American Heart Association and the European Society of Cardiology recommend intake of 1000 mg/d EPA+DHA to reduce CHD mortality.^{93,94} While lower doses may be sufficient (see Part 1), this population is at high risk for CHD death and most data to-date are from primary prevention studies. Thus, a target intake of 500-1000 mg/d (consistent with the largest secondary prevention trial to-date¹⁰) appears reasonable for individuals with established CHD. This level of intake can be achieved by consuming one 6-oz serving/week of fish richest in n-3 PUFAs (such as farmed salmon, anchovies, or herring), more frequently consuming other types of fish (Table 1), or by taking fish oil supplements. The maximum recommended dose of supplements without physician supervision is 3 g/d EPA+DHA.⁹⁵ Optimal levels of fish or fish oil intake for other clinical outcomes are not well-established.

For the general population (adolescents, men, and women not of childbearing age), there are no clear health effects of low level methylmercury exposure, and the strength of the evidence and the magnitude of the health benefits greatly exceeds the potential risk. Further investigation is needed to determine whether methylmercury exposure may modestly reduce the cardiovascular benefits of fish intake. For individuals concerned about mercury

exposure, consuming fish 1-2 servings/week and choosing a variety of different seafood will ensure low exposure, since most fish contain very low levels of mercury (Table 1). Individuals with very high fish consumption (≥ 5 servings/week) may choose to limit intake of the few species highest in mercury (Table 1).

DHA appears beneficial for early neurodevelopment. To minimize methylmercury exposure and possible sub-clinical neurologic harm, women of childbearing age and young children should avoid selected species: shark, swordfish, king mackerel, tilefish, and locally caught sportsfish per local advisories.^{26,45} However, emphasis must also be placed on adequate consumption of other fish and shellfish – up to 2 servings/wk, including up to 1 serving/wk of albacore tuna – to provide reasonable DHA^{26,45} and avoid further decreases in the already low consumption of seafood by women (74% of women of childbearing age and 85% of pregnant women consume <6 oz/week).^{96, 97}

Continued efforts to limit environmental contamination from organochlorine compounds are appropriate. However, the levels of PCBs and dioxins in fish are low, similar to levels in several other foods, and the associated possible risks are far smaller than the health benefits of fish intake. Thus, concern over PCBs/dioxins should not impact individual decisions regarding fish consumption (for locally caught freshwater fish, women of childbearing age should consult regional advisories).

In conclusion, the potential risks of fish intake must be considered in the context of the potential benefits. For major health outcomes among adults, the strength of the evidence and potential magnitudes of the benefits of modest fish intake (1-2 servings/week) greatly exceed those of the potential risks. For women of childbearing age, the benefits of modest fish intake, excepting a few selected species, also outweigh risks.

References

- Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: Evaluating the risks and the benefits. *JAMA* 2006;296:1855-99.
- Bang HO, Dyerberg J. Lipid metabolism and ischemic heart disease in Greenland Eskimos. In: Draper H, ed. *Advances in Nutrition Research*. New York, NY: Plenum Press;1980:1-22.
- Kromhout D, Bosschieter EB, de Lezenne Coulander C. The inverse relation between fish consumption and 20-year mortality from coronary heart disease. *N Engl J Med* 1985;312:1205-1209.
- Burr ML, Fehily AM, Gilbert JF, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;2:757-761.
- Dolecek TA, Granditis G. Dietary polyunsaturated fatty acids and mortality in the Multiple Risk Factor Intervention Trial (MRFIT). *World Rev Nutr Diet* 1991;66: 205-216.
- Siscovick DS, Raghunathan TE, King I, et al. Dietary intake and cell membrane levels of long-chain n-3 polyunsaturated fatty acids and the risk of primary cardiac arrest. *JAMA* 1995;274:1363-1367.
- Kromhout D, Feskens EJ, Bowles CH. The protective effect of a small amount of fish on coronary heart disease mortality in an elderly population. *Int J Epidemiol* 1995;24:340-345.
- Davignus ML, Stamler J, Orenica AJ, et al. Fish consumption and the 30-year risk of fatal myocardial infarction. *N Engl J Med* 1997;336:1046-1053.
- Albert CM, Hennekens CH, O'Donnell CJ, et al. Fish consumption and risk of sudden cardiac death. *JAMA* 1998;279:23-28.
- Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* 1999;354: 447-455.
- Oomen CM, Feskens EJ, Rasanen L, et al. Fish consumption and coronary heart disease mortality in Finland, Italy, and The Netherlands. *Am J Epidemiol* 2000; 151:999-1006.
- Yuan JM, Ross RK, Gao YT, Yu MC. Fish and shellfish consumption in relation to death from myocardial infarction among men in Shanghai, China. *Am J Epidemiol* 2001;154:809-816.
- Hu FB, Bronner L, Willett WC, et al. Fish and omega-3 fatty acid intake and risk of coronary heart disease in women. *JAMA* 2002;287:1815-1821.
- Albert CM, Campos H, Stampfer MJ, et al. Blood levels of long-chain n-3 fatty acids and the risk of sudden death. *N Engl J Med* 2002;346:1113-1118.
- Lemaitre RN, King IB, Mozaffarian D, Kuller LH, Tracy RP, Siscovick DS. n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* 2003; 77:319-325.
- Mozaffarian D, Lemaitre RN, Kuller LH, Burke GL, Tracy RP, Siscovick DS. Cardiac benefits of fish consumption may depend on the type of fish meal consumed: the Cardiovascular Health Study. *Circulation* 2003;107:1372-1377.
- Mozaffarian D, Ascherio A, Hu FB, et al. Interplay between different polyunsaturated fatty acids and risk of coronary heart disease in men. *Circulation* 2005; 111:157-164.
- Yokoyama M, Origasa H, Matsuzaki M, et al. Effects of eicosapentaenoic acid (EPA) on major cardiovascular events in hypercholesterolemic patients: The Japan EPA Lipid Intervention Study (JELIS). Paper presented at: American Heart Association Scientific Sessions; Nov 17, 2005; Dallas, TX.
- McLennan PL. Myocardial membrane fatty acids and the antiarrhythmic actions of dietary fish oil in animal models. *Lipids* 2001;36 Suppl:S111-114.
- Leaf A, Kang JX, Xiao YF, Billman GE. Clinical prevention of sudden cardiac death by n-3 polyunsaturated fatty acids and mechanism of prevention of arrhythmias by n-3 fish oils. *Circulation* 2003;107:2646-2652.
- Wang C, Harris WS, Chung M, et al. n-3 Fatty acids from fish or fish-oil supplements, but not (alpha)-linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review. *Am J Clin Nutr* 2006;84:5-17.
- U.S. Environmental Protection Agency. Mercury Study Report to Congress. Available at: <http://www.epa.gov/mercury/report.htm>. Accessed January 24, 2006.
- U.S. Geological Survey. Mercury in the Environment. Available at: <http://www.usgs.gov/themes/factsheet/146-00/>. Accessed October 25, 2005.
- Committee on the Toxicological Effects of Methylmercury; Board on Environmental Studies and Toxicology; Commission on Life Sciences; National Research Council. *Toxicological Effects of Methylmercury*. Washington, DC: National Academy Press; 2000.
- The Risk Assessment Information System. Toxicity Summary for Mercury. Available at: http://risk.lsd.ornl.gov/tox/profiles/mercury_f_V1.shtml. Accessed January 24, 2006.
- U.S. Food And Drug Administration. Center for Food Safety and Applied Nutrition. Seafood Information and Resources. Available at: <http://www.cfsan.fda.gov/seafood1.html>. Accessed January 30, 2006.
- World Health Organization. Assessment of the health risk of dioxins: re-evaluation of the Tolerable Daily Intake (TDI): WHO Consultation, May 25-29, Geneva, Switzerland; 1998.
- U.S. Environmental Protection Agency. National Center for Environmental Assessment. Dioxin and Related Compounds. Available at: <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=55264>. Accessed March 14, 2006.
- U.S. Environmental Protection Agency. Polychlorinated Biphenyls (PCBs). Available at: <http://www.epa.gov/opptintr/pcb/>. Accessed March 14, 2006.
- Verbeke W, Sioen I, Pieniak Z, Van Camp J, De Henauw S. Consumer perception versus scientific evidence about health benefits and safety risks from fish consumption. *Public Health Nutr* 2005;8:422-429.
- Center for Food Nutrition and Agriculture Policy. University of Maryland. Real Mercury Facts. Available at: <http://www.realmercuryfacts.org/index.htm>. Accessed March 23, 2006.
- Gochfeld M. Cases of mercury exposure, bioavailability, and absorption. *Ecotoxicol Environ Saf* 2003;56:174-179.
- U.S. Environmental Protection Agency. Methylmercury (MeHg) (CASRN 22967-92-6). Available at: <http://www.epa.gov/iris/subst/0073.htm>. Accessed May 1, 2006.
- McDowell MA, Dillon CF, Osterloh J, et al. Hair mercury levels in U.S. children and women of childbearing age: reference range data from NHANES 1999-2000. *Environ Health Perspect* 2004;112:1165-1171.
- Grandjean P, Weihe P, White RF, et al. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 1997;19:417-428.
- Grandjean P, Weihe P, White RF, Debes F. Cognitive performance of children prenatally exposed to "safe" levels of methylmercury. *Environ Res* 1998;77:165-172.
- Kjellstrom T. Physical and mental development of children with prenatal exposure to mercury from fish. Stage II: interviews and psychological tests at age 6. Stockholm, Sweden: National Swedish Environmental Protection Board; 1989;Report 3642. Solna, Sweden:National Swedish Environmental Protection Board.
- Crump KS, Kjellstrom T, Shipp AM, Silvers A, Stewart A. Influence of prenatal mercury exposure upon scholastic and psychological test performance: benchmark analysis of a New Zealand cohort. *Risk Anal* 1998;18:701-713.
- Jedrychowski W, Jankowski J, Flak E, et al. Effects of prenatal exposure to mercury on cognitive and psychomotor function in one-year-old infants: Epidemiologic Cohort Study in Poland. *Ann Epidemiol* 2006;16(6):439-47.
- Palumbo DR, Cox C, Davidson PW, et al. Association between prenatal exposure to methylmercury and cognitive functioning in Seychellois children: a reanalysis of the McCarthy Scales of Children's Ability from the main cohort study. *Environ Res* 2000;84:81-88.
- Davidson PW, Palumbo D, Myers GJ, et al. Neurodevelopmental outcomes of Seychellois children from the pilot cohort at 108 months following prenatal exposure to methylmercury from a maternal fish diet. *Environ Res* 2000;84:1-11.
- Oken E, Wright RO, Kleinman KP, et al. Maternal fish consumption, hair mercury, and infant cognition in a U.S. Cohort. *Environ Health Perspect* 2005;113:1376-1380.
- Daniels JL, Longnecker MP, Rowland AS, Golding J. Fish intake during pregnancy and early cognitive development of offspring. *Epidemiol* 2004;15:394-402.
- Spurgeon A. Prenatal methylmercury exposure and developmental outcomes: review of the evidence and discussion of future directions. *Environ Health Perspect* 2006;114:307-312.
- U.S. Environmental Protection Agency. What you need to know about mercury in fish and shellfish. Available at: <http://www.epa.gov/waterscience/fishadvice/advice.html>. Accessed March 25, 2006.
- Rice DC. The US EPA reference dose for methylmercury: sources of uncertainty. *Environ Res* 2004;95:406-413.
- Ahlqvist M, Bengtsson C, Lapidus L, Gergdahl IA, Schutz A. Serum mercury concentration in relation to survival, symptoms, and diseases: results from the prospective population study of women in Gothenburg, Sweden. *Acta Odontol Scand* 1999; 57:168-174.
- Hallgren CG, Hallmans G, Jansson JH, et al. Markers of high fish intake are associated with decreased risk of a first myocardial infarction. *Br J Nutr* 2001;86:397-404.
- Gualler E, Sanz-Gallardo MI, van't Veer P, et al. Mercury, fish oils, and the risk of myocardial infarction. *N Engl J Med* 2002;347:1747-1754.
- Yoshizawa K, Rimm EB, Morris JS, et al. Mercury and the risk of coronary heart disease in men. *N Engl J Med* 2002;347:1755-1760.

51. Virtanen JK, Voutilainen S, Rissanen TH, et al. Mercury, fish oils, and risk of acute coronary events and cardiovascular disease, coronary heart disease, and all-cause mortality in men in eastern Finland. *Arterioscler Thromb Vasc Biol* 2005; 25:228-233.
52. Rissanen T, Voutilainen S, Nyyssonen K, Lakka TA, Salonen JT. Fish oil-derived fatty acids, docosahexaenoic acid and docosapentaenoic acid, and the risk of acute coronary events: the Kuopio ischaemic heart disease risk factor study. *Circulation* 2000;102:2677-2679.
53. Risher JF, Murray HE, Prince GR. Organic mercury compounds: human exposure and its relevance to public health. *Toxicol Ind Health* 2002;18:109-160.
54. Risher JF. Too much of a good thing (fish): methylmercury case study. *J Environ Health* 2004;67:9-14, 28.
55. Lebel J, Mergler D, Branches F, et al. Neurotoxic effects of low-level methylmercury contamination in the Amazonian Basin. *Environ Res* 1998;79:20-32.
56. Yokoo EM, Valente JG, Grattan L, Schmidt SL, Platt I, Silbergeld EK. Low level methylmercury exposure affects neuropsychological function in adults. *Environ Health* 2003;2:8.
57. Auger N, Kofman O, Kosatsky T, Armstrong B. Low-level methylmercury exposure as a risk factor for neurologic abnormalities in adults. *Neurotoxicology* 2005;26:149-157.
58. Weil M, Bressler J, Parsons P, Bolla K, Glass T, Schwartz B. Blood mercury levels and neurobehavioral function. *JAMA* 2005;293:1875-1882.
59. Johansson N, Basun H, Winblad B, Nordberg M. Relationship between mercury concentration in blood, cognitive performance, and blood pressure, in an elderly urban population. *Biometals* 2002;15:189-195.
60. He K, Song Y, Daviglius ML, et al. Fish consumption and incidence of stroke: a meta-analysis of cohort studies. *Stroke* 2004;35:1538-1542.
61. Morris MC, Evans DA, Tangney CC, Bienias JL, Wilson RS. Fish consumption and cognitive decline with age in a large community study. *Arch Neurol* 2005;62:1849-1853.
62. Peet M, Stokes C. Omega-3 fatty acids in the treatment of psychiatric disorders. *Drugs* 2005;65:1051-1059.
63. Young G, Conquer J. Omega-3 fatty acids and neuropsychiatric disorders. *Reprod Nutr Dev* 2005;45:1-28.
64. U.S. Food And Drug Administration. Food and Drug Administration Total Diet Study. Available at: <http://vm.cfsan.fda.gov/~comm/ds-toc.html>. Accessed February 1, 2006.
65. U.S. Department of Agriculture. Food Safety and Inspection Service. Dioxins and Dioxin-Like Compounds In the U.S. Domestic Meat and Poultry Supply. Available at: http://www.fsis.usda.gov/PDF/Dioxin_Report_0605.pdf. Accessed March 24, 2006.
66. Liem AK, Furst P, Rappe C. Exposure of populations to dioxins and related compounds. *Food Addit Contam* 2000;17:241-259.
67. Gomara B, Bordajandi LR, Fernandez MA, et al. Levels and trends of polychlorinated dibenzo-p-dioxins/furans (PCDD/Fs) and dioxin-like polychlorinated biphenyls (PCBs) in Spanish commercial fish and shellfish products, 1995-2003. *J Agric Food Chem* 2005; 53:8406-8413.
68. National Center for Environmental Assessment, U.S. Environmental Protection Agency. PCBs: Cancer dose-response assessment and application to environmental mixtures: Washington, D.C.; 1996.
69. Schechter A, Cramer P, Boggess K, et al. Intake of dioxins and related compounds from food in the U.S. population. *J Toxicol Environ Health A* 2001;63:1-18.
70. Hites RA, Foran JA, Carpenter DO, Hamilton MC, Knuth BA, Schwager SJ. Global assessment of organic contaminants in farmed salmon. *Science* 2004;303:226-229.
71. Hamilton MC, Hites RA, Schwager SJ, Foran JA, Knuth BA, Carpenter DO. Lipid composition and contaminants in farmed and wild salmon. *Environ Sci Technol* 2005;39: 8622-8629.
72. Foran JA, Good DH, Carpenter DO, Hamilton MC, Knuth BA, Schwager SJ. Quantitative analysis of the benefits and risks of consuming farmed and wild salmon. *J Nutr* 2005;135:2639-2643.
73. U.S. Environmental Protection Agency. Guidance for Assessing Chemical Contaminant Data for Use In Fish Advisories. Volume 2: Risk Assessment and Fish Consumption Limits – Third Edition. Washington, D.C.; 2003.
74. MacLean CH, Newberry SJ, Mojica WA, et al. Effects of omega-3 fatty acids on cancer risk: a systematic review. *JAMA* 2006;295:403-415.
75. Mori TA, Beilin LJ. Omega-3 fatty acids and inflammation. *Curr Atheroscler Rep* 2004; 6:461-467.
76. Mickleborough TD, Lindley MR, Ionescu AA, Fly AD. Protective effect of fish oil supplementation on exercise-induced bronchoconstriction in asthma. *Chest* 2006;129:39-49.
77. Jacobson JL, Jacobson SW. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 1996;335:783-789.
78. Patandin S, Lanting CI, Mulder PG, Boersma ER, Sauer PJ, Weisglas-Kuperus N. Effects of environmental exposure to polychlorinated biphenyls and dioxins on cognitive abilities in Dutch children at 42 months of age. *J Pediatr* 1999;134:33-41.
79. Grandjean P, Weihe P, Burse VW, et al. Neurobehavioral deficits associated with PCB in 7-year-old children prenatally exposed to seafood neurotoxicants. *Neurotoxicol Teratol* 2001;23:305-317.
80. Ribas-Fito N, Sala M, Kogevinas M, Sunyer J. Polychlorinated biphenyls (PCBs) and neurological development in children: a systematic review. *J Epidemiol Community Health* 2001;55:537-546.
81. Stewart PW, Reihman J, Lonky EI, Darvill TJ, Pagano J. Cognitive development in preschool children prenatally exposed to PCBs and MeHg. *Neurotoxicol Teratol* 2003; 25:11-22.
82. Schantz SL, Widholm JJ, Rice DC. Effects of PCB exposure on neuropsychological function in children. *Environ Health Perspect* 2003;111:357-576.
83. Nakajima S, Saijo Y, Kato S, et al. Effects of prenatal exposure to polychlorinated biphenyls and dioxins on mental and motor development in Japanese children at 6 months of age. *Environ Health Perspect* 2006;114:773-778.
84. Daniels JL, Longnecker MP, Klebanoff MA, et al. Prenatal exposure to low-level polychlorinated biphenyls in relation to mental and motor development at 8 months. *Am J Epidemiol* 2003;157:485-492.
85. Gray KA, Klebanoff MA, Brock JW, et al. In utero exposure to background levels of polychlorinated biphenyls and cognitive functioning among school-age children. *Am J Epidemiol* 2005;162:17-26.
86. Judd N, Griffith WC, Faustman EM. Contribution of PCB exposure from fish consumption to total dioxin-like dietary exposure. *Regul Toxicol Pharmacol* 2004;40:125-135.
87. Jacobs MN, Covaci A, Schepens P. Investigation of selected persistent organic pollutants in farmed Atlantic salmon (*Salmo salar*), salmon aquaculture feed, and fish oil components of the feed. *Environ Sci Technol* 2002;36:2797-2805.
88. Storelli MM, Storelli A, Marcotrigiano GO. Polychlorinated biphenyls, hexachlorobenzene, hexachlorocyclohexane isomers, and pesticide organochlorine residues in cod-liver oil dietary supplements. *J Food Prot* 2004;67:1787-1791.
89. Jimenez B, Wright C, Kelly M, Startin JR. Levels of PCDDs, PCDFs and non-ortho PCBs in dietary supplement fish oil obtained in Spain. *Chemosphere* 1996;32:461-467.
90. Food Safety Authority of Ireland. Summary of investigation of dioxins, furans, and PCBs in farmed salmon, wild salmon, farmed trout and fish oil capsules, March 2002. Available at: http://www.fsai.ie/surveillance/food/surveillance_food_summarydioxins.asp. Accessed March 31, 2006.
91. Foran SE, Flood JG, Lewandrowski KB. Measurement of mercury levels in concentrated over-the-counter fish oil preparations: is fish oil healthier than fish? *Arch Pathol Lab Med* 2003;127:1603-1605.
92. Brown AJ, Pang E, Roberts DC. Persistent changes in the fatty acid composition of erythrocyte membranes after moderate intake of n-3 polyunsaturated fatty acids: study design implications. *Am J Clin Nutr* 1991;54:668-673.
93. Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* 2002;106:2747-2757.
94. Van de Werf F, Ardissino D, Betriu A, et al. Management of acute myocardial infarction in patients presenting with ST-segment elevation. The Task Force on the Management of Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2003;24:28-66.
95. U.S. Food And Drug Administration. Substances Affirmed as Generally Recognized as Safe: Menhaden Oil. Available at: <http://a257.g.akamaitech.net/1/257/2422/14mar2001/0800/edocket.access.gpo.gov/2004/04-811.htm>. Accessed April 5, 2006.
96. Schober SE, Sinks TH, Jones RL, et al. Blood mercury levels in US children and women of childbearing age, 1999-2000. *JAMA* 2003;289:1667-1674.
97. Oken E, Kleinman KP, Berland WE, Simon SR, Rich-Edwards JW, Gillman MW. Decline in fish consumption among pregnant women after a national mercury advisory. *Obstet Gynecol* 2003;102:346-351.
98. Thannum J, Great Lakes Indian Fish & Wildlife Commission. Tribally sold Lake Superior fish easily meet FDA restrictions for chemical contaminants. Available at: http://www.glifwc.org/pub/summer00/fish_contaminants.htm. Accessed March 25, 2006.



Dr. Dariush Mozaffarian obtained his undergraduate degree from Stanford University and his MD from Columbia College of Physicians and Surgeons. He completed his Internal Medicine residency at Stanford University and his Cardiology fellowship at the University of Washington. He also obtained a Masters in Public Health from the University of Washington and a Doctorate in Epidemiology from Harvard. He holds appointments in the Department of Medicine at Harvard Medical School and in the Department of Epidemiology at the Harvard School of Public Health. His primary research interests are the effects of dietary habits on cardiovascular health and disease, including coronary heart disease, sudden death, heart failure, and atrial fibrillation. He has authored numerous original publications on these topics. He is particularly interested in the effects of dietary fats, especially omega-3 fatty acids (found in seafood and plant sources) and trans fatty acids (found in partially hydrogenated oils). He has also elucidated cardiovascular effects of other dietary factors, such as dietary fiber, refined carbohydrates, and saturated fats. His research efforts are funded by multiple grants from the National Institutes of Health.

Disclosure: Dr. Dariush Mozaffarian has no conflicts of interest in association with the contents of this issue.

Brigham and Women's Hospital,
Cardiovascular Division website:
www.heartdoc.org

This publication is made possible by an educational grant from
Novartis Pharmaceuticals Corporation

© 2006 Brigham and Women's Hospital, Boston, Massachusetts, which is solely responsible for the contents. The opinions expressed in this publication do not necessarily reflect those of the publisher or sponsor, but rather are those of the author based on the available scientific literature. Publisher: **SNELL Medical Communication Inc.** in cooperation with Brigham and Women's Hospital, Boston, Massachusetts. *Cardiology Rounds* is a registered trade mark of SNELL Medical Communication Inc. All rights reserved. The administration of any therapies discussed or referred to in *Cardiology Rounds* should always be consistent with the recognized prescribing information as required by the FDA. **SNELL Medical Communication Inc.** is committed to the development of superior Continuing Medical Education.