

**Harvard Medical School Department of  
Continuing Education and the Cardiovascular  
Division of the Department of Medicine,  
Brigham and Women's Hospital**



***Cardiology Rounds***  
**February 2004**

**Prediction of Coronary Heart Disease Events.  
Part 2: The Contribution of Lifestyle Factors and New Issues.**

By Peter W. F. Wilson, M.D.

**Objectives:**

In this second component of *Cardiology Rounds*, Dr. Wilson discusses newer potential risk factors and how to ascertain whether they offer incremental predictive value. The reader should obtain a better understanding of the multi-factorial aspects of individual risk factor assessment and be cognizant of new candidates.

**TEST:**

1. Although observational studies indicate that greater intake of vitamins B, C, and E have been associated with lower cardiovascular risks, randomized controlled clinical trials have yet to demonstrate that dietary supplementation of these vitamins reduces cardiovascular risk.  
True       False
2. Moderate alcohol intake (less than two drinks per day in men and less than one drink per day in women) reduces cardiovascular risk by lowering triglycerides and LDL cholesterol.  
True       False
3. Physical fitness, whether assessed by the degree of leisure-time physical activity or longer exercise treadmill times, is associated with reduced risk for experiencing cardiovascular disease in men but not women.  
True       False
4. The prevalence of obesity in the United States defined as a body mass index (BMI)  $\geq 30$  kg/meter<sup>2</sup> has more than doubled in the past 30 years.  
True       False
5. Measurements of inflammatory markers, specifically high sensitivity C-reactive protein (hsCRP) provides incremental adjunctive information for cardiovascular risk factor assessment.  
True       False

6. The most important issues for consideration of newer risk factors is
- a. accuracy
  - b. cost
  - c. variability
  - d. incremental utility of the test
7. Discrimination has the following characteristics.
- a. Ability to validate the predictive capability of a diagnostic approach in one study and apply it to another study
  - b. Ability to identify persons who will develop disease and those who will not, derived from the sensitivity and specificity
  - c. The likelihood of developing disease when the prevalence is 20%
  - d. A ratio derived from the standard deviation and mean for a test
8. The key risk for a patient to know is:
- a. relative risk
  - b. excess risk compared to others in the population
  - c. absolute risk
  - d. population attributable risk
9. Arterial calcification in African Americans for CHD prediction may have special limitations because
- a. left ventricular hypertrophy is much more common
  - b. familial hypercholesterolemia is more prevalent
  - c. dyslipidemia is more frequent
  - d. calcification is less prevalent in African Americans

To receive AMA category 1 credit, you must correctly answer 60% of the test questions.

Harvard Medical School is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

Harvard Medical School designates this educational activity for a maximum of 1 category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those credits that he/she actually spent in the educational activity.

This program was issued in February 2004. All tests must be returned by June 30, 2004.

Please send the completed test and a check for \$25 US. (Check made payable to: Harvard Medical School) to: Harvard Medical School/Department of Continuing Education, Box 825, Boston, MA 02117-825.

We will return your corrected test and a certificate upon successful completion.

Name \_\_\_\_\_

Address \_\_\_\_\_

Phone \_\_\_\_\_

Email \_\_\_\_\_

Fax \_\_\_\_\_