

**Harvard Medical School Department of
Continuing Education and the Cardiovascular
Division of the Department of Medicine,
Brigham and Women's Hospital**



Cardiology Rounds
December 2005

Diastolic and Systolic Heart Failure – Similarities and Differences. Part 2.

By Kanu Chatterjee, MB, FRCP, FCCP, FACC, MACP

Objectives:

In the second of a two-part series that compares and contrasts the morphologic and functional characteristics of patients with diastolic and systolic heart failure, the major emphases are on structural remodeling and therapeutic options. This issue reviews the therapeutic strategies for the treatment of these disorders and underscores their uneven progress, highlighting the need for further research. This issue of *Cardiology Rounds* will help readers acquire a better understanding of the morphologic and functional differences of the left ventricle that characterize these two forms of heart failure.

Questions:

1. Patients with diastolic heart failure have derangements in relaxation and passive stiffness that result in an increased left ventricular filling pressure relative to the ventricular volume.

True False

2. Left ventricular morphologic and functional characteristics of patients with diastolic heart failure commonly include all the following except:

- a. increased wall thickness
- b. reduced ventricular volumes
- c. increased wall stress
- d. increased mass
- e. increased ejection fraction

3. Elevations in pulmonary venous pressure with associated increased pulmonary vascular resistance is a common feature in both systolic and diastolic heart failure.

True False

4. Diuretics are the principal therapeutic agents to relieve congestive symptoms in both systolic and diastolic heart failure patients.

True False

5. Angiotensin-converting enzyme (ACE) inhibitors have been demonstrated to improve mortality in patients with both systolic and diastolic heart failure.

True False

6. Mechanical devices such as cardiac resynchronization therapy and implantable defibrillators improve prognosis of both patients with systolic and diastolic heart failure.

True False

7. Cardiac transplantation should not be considered as a therapeutic option for patients with severe refractory diastolic heart failure.

True False

To receive AMA category 1 credit, you must correctly answer 60% of the test questions.

Harvard Medical School is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

Harvard Medical School designates this educational activity for a maximum of 1 category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those credits that he/she actually spent in the educational activity.

This program was issued in December 2005. All tests must be returned by April 30, 2006.

Please send the completed test and a check for \$25 US. (Check made payable to: Harvard Medical School) to: Harvard Medical School/Department of Continuing Education, Box 825, Boston, MA 02117-825.

Please keep a copy of your test before submission. A certificate will be sent upon successful completion of the test, along with the answer key, after the deadline date, as indicated.

Name _____

Address _____

Phone _____

Email _____

Fax _____